

DESCRIPTIONDRIED BLOOD FACTOR COMPOSITION COMPRISING TREHALOSE5 Cross-Reference to Related Applications

This application is a continuation application of co-pending application U.S. Serial No. 09/888,734, filed June 25, 2001; which is a continuation application of U.S. Serial No. 08/875,796, filed October 30, 1998, now U.S. Patent No. 6,649,386, which was filed as a National Stage Application of International Application Number
10 PCT/GB96/00119, filed January 19, 1996, published, pursuant to PCT Article 21(2); which claims priority to Great Britain Application GB 9501040.1, January 19, 1995.

Background of the Invention

 This invention relates to dried compositions of blood factors for reconstitution
15 with water or aqueous solutions.

 Blood factors, particularly factor VIII and factor IX, are now the standard treatment for diseases caused by a lack of the appropriate factor, in particular haemophilia. The blood factor has generally been derived from human blood by various
20 extraction techniques, for example as disclosed in EP-A-0083483, or by expression in genetically modified microorganisms, for example as disclosed in EP-A-0160457 and EP-A-0182448.

 Blood factor products such as factor VIII are highly delicate, unstable proteins.
25 They are usually supplied in the form of frozen solutions in an appropriate buffer or, more generally, as freeze-dried powders. Even the freeze-dried powders must be kept cold during storage. In order to stabilise the freeze-dried material, commercial products contain a stabilising protein, in particular human serum albumin (HSA). It has not been thought possible to prepare a dry blood factor composition which is stable at ambient
30 temperatures and at pasteurisation temperatures (e.g. 60°C) in the absence of HSA.

However, the presence of HSA introduces considerable problems of purification since it is essential that the protein is free of viral contamination. The use of recombinant HSA to overcome these problems is expensive.

5 Trehalose is known to be a highly effective stabilising agent for delicate proteins, as disclosed in US-A-4891319, enabling proteins to be dried at temperatures above freezing. We have now found that if trehalose is used to stabilise a blood factor product, not only can the product be dried with or without freezing, but also the product is stable even when retained at a temperature of 60°C for an extended period, in the complete
10 absence of HSA. According to the present invention therefore we provide a stable dried blood factor composition containing a stabilising amount of trehalose in the absence of albumin.

15 In general, any stabilising amount of trehalose may be used and an excess in general causes no problems. Indeed, the presence of trehalose aids the rehydration process and is physiologically acceptable for injection, being rapidly metabolised to glucose. The composition is particularly suited to formulations of factor VIII, which may also contain appropriate buffering and ion-reinforcing salts, in particular a source of calcium. In general, a ratio of about 1.0 to 1.5 mg of calcium ions per unit of factor VIII
20 is appropriate.

25 Other buffering and modifying agents may also be present in the dried material for reconstitution to the injection solution, for example histidine. However, we have found that the level of salts, particularly sodium chloride, present can affect the preservation on drying. It is thought desirable for the commercial product for injection to have an isotonic salt concentration. However, the processing formulations which are freeze-dried are desirably hypertonic, typically containing about 500 mM NaCl (isotonic NaCl = 150 mM), as this is considered to help stabilise the blood factor. As a result, commercial freeze-dried formulations generally have a high salt content and are

reconstituted for injection with the appropriate amount of sterile water to obtain an isotonic solution.

5 A considerably reduced salt content is preferred for the dried material of the invention and, in general a solution of about 500 units of Factor VIII per ml to be dried should preferably contain less than 200 mM e.g. 75 to 150 mM, NaCl, especially about 100 mM, or even lower, e.g. 20 to 50 mM, especially about 22 to 30 mM. Low salt preparations possess a higher dry stability. The dried product can be reconstituted to the desired salt level with a saline solution instead of the conventional water. In general, the
10 molar ratio of trehalose to salt should be above 1:1, especially above 2.5:1 e.g. above 10:1, preferably above 12.5:1.

The dried composition may be obtained by drying an appropriate solution of the blood factor containing the correct proportions of trehalose and other desired
15 components. In general, the solution that is dried should simply contain all the components required in the reconstituted injection solution, although the solution for drying may not necessarily be at the same dilution. Typically, the solution for drying will contain from 1 to 1000 units of factor VIII per ml. The methods of drying may include freeze drying, vacuum drying and spray-drying. A particularly preferred method
20 according to the invention comprises vacuum drying at a temperature no greater than 25°C, preferably no greater than 10°C, to form a foam, thus maximising the exposed surface and the drying effect.

The following examples illustrate the invention further.

25 Example 1

Recombinant factor VIII was received as a deep frozen solution containing approximately 2000 to 2500 units/ml in the manufacturer's high salt buffer. The thawed solution was dialysed against a buffer solution containing 500 mM NaCl, 15 mM CaCl₂
30 and 10 mM histidine at pH 6.8. The dialysed protein was diluted in the same buffer, but

with added trehalose to give a final concentration of 500 units per ml and 10% by weight trehalose at pH 6.8. This solution was vacuum dried in 1 ml aliquots. Vacuum was reduced step-wise from atmospheric to 4 Pa (30 mTorr) to avoid freezing the sample. The temperature of the sample was not allowed to rise above 12°C until the formation of a foam, after which the temperature was kept below 30°C. Total drying times were 24 to 28 hours.

The samples were stored for 0, 1.5, 3 and 6 months at 40°C and then reconstituted in 5 ml aliquots of sterile distilled water before being tested for activity. The results are shown in the following table in comparison with a commercial freeze-dried product containing HSA. Both the test and commercial samples have a high salt content. The post-drying results show that with trehalose it is possible to dry factor VIII successfully in the absence of HSA, but that a high salt content is unsatisfactory for long term storage, even in the presence of HSA.

Percentage of initial activity

Time (months)	Sample	Commercial Product
0	100.0	100.0
1.5	86.8	95.3
3	75.1	71.2
6	76.6	63.6

Example 2

Samples were dried as described in Example 1 but using a buffer formulation comprising 100 mM NaCl, 15 mM CaCl₂, 15mM histidine and 1.27 molar trehalose (48%) and stored at 60°C before reconstitution. The results are given in the following table, in which the activity is measured on an ACL 100 automated coagulometer (Instrumentation Laboratory SpA, Milan, Italy). The test sample, with a low salt content showed no significant loss of activity on storage, even after four weeks of 60°C.

Percentage of initial activity recovered

Wet control	100.0
Post-drying	95.5
Two weeks	96.0
Four weeks	96.8

Example 3

Two formulations were prepared containing different salt concentrations as shown:

5

Formulation A		Formulation B	
NaCl	0.13%	NaCl	1.03%
CaCl ₂	0.011%	CaCl ₂	0.011%
L-histidine	0.12%	L-histidine	0.12%
Tris	0.002%	Tris	0.002%
Tween 80	0.002%	Tween 80	0.002%
PEG 3350	0.004%	PEG 3350	0.004%
Trehalose	7.5%	Trehalose	7.5%
Factor VIII	50 U/ml	Factor VIII	50 U/ml
Water to 100%		Water to 100%	

10

15

10ml portions of the formulations were dispensed into separate 30ml vials so as to give a concentration of factor VIII of 500 units/vial.

20

Freeze-drying was performed in a Laconco (Lyph-lock 12 stoppering) freeze drier. Initially, the samples were cooled to -40°C and then placed under vacuum, before being warmed to -35°C. After 80 hours, the samples were warmed at a rate of 2.5°C/h until the shelf temperature reached 25°C. The samples were then kept at 25°C for two hours before being sealed under vacuum and removed from the drier.

25

After drying, the samples were rehydrated with 10ml of water and the concentration of factor VIII was measured twice (Assays 1 and 2). The results are given in the following table and are expressed as a percentage of the concentration of Factor VIII with respect to the prefill control (which had been frozen at -70°C). From the results shown, it can be concluded that Factor VIII can be successfully freeze dried in a trehalose based formulation in the absence of HSA.

Sample	Assay 1	Assay 2
Formulation A	77%	85%
Formulation B	91%	94%

Abstract of the Disclosure

A stable blood factor composition contains a stabilising amount of trehalose in the absence of human serum albumin to provide a product stable at up to 60°C.